

# Weak levitation of 2D delocalized states in a magnetic field

T. V. Shahbazyan and M. E. Raikh

*Department of Physics, University of Utah, Salt Lake City, UT 84112*

## Abstract

The deviation of the energy position of a delocalized state from the center of Landau level is studied in the framework of the Chalker-Coddington model. It is demonstrated that introducing a weak Landau level mixing results in a shift of the delocalized state up in energy. The mechanism of a levitation is a neighboring - Landau level - assisted resonant tunneling which “shunts” the saddle-points. The magnitude of levitation is shown to be independent of the Landau level number.

PACS numbers: 73.20.Jc, 73.30.+h, 73.40.Hm

It is commonly accepted that electronic spectrum of a two-dimensional disordered system in a strong magnetic field contains only a single delocalized state per Landau level (LL). First conjectured by Halperin [1], this conclusion was later drawn from the consideration based on scaling ideas [2,3] and confirmed by numerical simulations in which the divergence of the localization length with energy approaching to the position of the delocalized state was studied [4–6].

To compromise this conjecture with the absence of delocalized states at zero field ( $B = 0$ ) Khmelnitskii [7] has suggested that with decreasing  $B$  the energy positions of delocalized states depart gradually from the centers of Landau levels and float upwards to reach an infinity as  $B \rightarrow 0$ . In order to obtain the positions of delocalized states,  $E_n(B)$ , Khmelnitskii has pointed out that the system of scaling equations [2] for conductivities,  $\sigma_{xx}$ ,  $\sigma_{xy}$  should be solved together with the boundary conditions that at small scales those are given by their classical expressions:  $\sigma_{xx}^{cl} = \sigma_0(1 + \omega_c^2\tau^2)^{-1}$ ,  $\sigma_{xy}^{cl} = \sigma_0\omega_c\tau(1 + \omega_c^2\tau^2)^{-1}$ , where  $\sigma_0$  is the Drude conductivity (proportional to the Fermi energy  $E_F$ ),  $\omega_c$  is the cyclotron frequency and  $\tau$  is the elastic scattering time. Since the scaling equations are periodic in  $\sigma_{xy}$ , it follows that  $\sigma_{xx}(E_F)$  remains finite at large scales each time when  $\sigma_{xy}^{cl} = (n + 1/2)e^2/2\pi\hbar$ . This gives  $E_n = \hbar\omega_c(n + 1/2)[1 + (\omega_c\tau)^{-2}]$ . In support of the above picture Laughlin [8] has presented an argument that the delocalized states cannot disappear discontinuously at any finite magnetic field.

The levitation of the delocalized states according to [7,8] is sketched in Fig. 1. The above dependence  $E_n(B)$ , treated as a dependence of  $\rho_{xx}^0 = \sigma_0^{-1}$  versus the inverse filling factor is, in fact, an essential constituent of the global phase diagram of the quantum Hall effect (QHE) introduced by Kivelson, Lee, and Zhang [9] (namely, an integer QHE part of this diagram). Recent experiments [10–13], in which insulator - quantum Hall conductor - insulator transition was observed with increasing magnetic field, were interpreted in terms of the global phase diagram [9]. Using Fig. 1 the interpretation presented in [10,12,13] can be reformulated as follows. If the Fermi level at zero field lies within some certain range, as shown in Fig. 1, then the horizontal line,  $E_F(B) = \text{const}$ , crosses the curve  $E_0(B)$  at two

different values of  $B$ . This implies that for  $B$  in between these values the system is in the QHE regime while for smaller or larger  $B$  the system is in the insulating phase. The latest experiment [14], carried out at very low temperatures, allowed, in fact, a direct measurement of  $E_0(B)$  dependence by tracing the positions of peaks in  $\sigma_{xx}$ , measured as a function of gate voltage, at different magnetic fields. The measured dependence  $E_0(B)$  exhibits a minimum in the region of  $B$  where spin levels are not resolved.

Numerical simulations [15] for the case of a short-range disorder indeed indicated an upward (from  $E = \hbar\omega_c/2$ ) shift of the energy position of the extended state when three lowest LL were taken into account. However, no microscopic theory for such a levitation has been developed so far. In the present paper we point out that such a theory can be developed in the region where the departure of the delocalized state from the center of the Landau band is relatively small. We demonstrate that the microscopic mechanism for the levitation is the resonant scattering of an electron with energy near the center of LL by localized states from neighboring LL.

We consider the case of a smooth random potential and adopt the network model by Chalker and Coddington [16] which we generalize in order to study the effects of neighboring LL. In this model delocalization results from a quantum tunneling of an electron through saddle-points of a smooth potential which are connected by equipotential lines. The randomness of the potential is included by assuming the phase acquired by electron traversing a link (equipotential line) to be random. The probability amplitudes of incoming waves  $Z_1$  and  $Z_2$  and outgoing waves  $Z_3$  and  $Z_4$  (see Fig. 2) are related as

$$\begin{pmatrix} Z_1 \\ Z_3 \end{pmatrix} = \mathbf{M} \begin{pmatrix} Z_4 \\ Z_2 \end{pmatrix}, \quad (1)$$

where the matrix  $\mathbf{M}$  can be decomposed into the product

$$\mathbf{M} = \begin{pmatrix} e^{i\phi_1} & 0 \\ 0 & e^{i\phi_2} \end{pmatrix} \begin{pmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{pmatrix} \begin{pmatrix} e^{i\phi_3} & 0 \\ 0 & e^{i\phi_4} \end{pmatrix}, \quad (2)$$

with parameter  $\theta$  characterizing the tunneling and  $\phi_i$  being gauge phases. For the potential

expanded near the saddle point,  $V(x, y) = V_0 - m\Omega_x^2 x^2/2 + m\Omega_y^2 y^2/2$ ,  $m$  being the electron mass, this parameter can be presented as [18]

$$\sinh \theta = \exp\left(\frac{E - V_0}{\gamma}\right), \quad \gamma = \hbar\Omega_x\Omega_y/\pi\omega_c, \quad (3)$$

where  $E$  is energy of electron measured from the center of LL. Since the saddle-point heights,  $V_0$ , are distributed *symmetrically* around the value  $V_0 = 0$  [corresponding to  $\theta = \theta_c \equiv \ln(1 + \sqrt{2})$ ], the delocalized states occurs at zero energy (Chalker and Coddington assumed  $V_0 = 0$  at all saddle points; later simulations [17] with randomly distributed  $V_0$  demonstrated the universality of the critical exponent).

Assume now that the energy is close to the center of  $n = 0$  LL and study the change in the structure of electron states caused by  $n = 1$  LL. The relevant  $n = 1$  LL states are those with energies close to the center of  $n = 0$  LL. The equipotentials corresponding to these states are depicted schematically in Fig. 2 by dashed loops. The prime role of these equipotentials is to shift the maximum of the density of states up from  $E = 0$ . However, such a shift does not affect the position of delocalized state unless a coupling between the equipotentials of  $n = 0$  and  $n = 1$  LL is introduced. This coupling is illustrated in Fig. 2a. The line connecting the equipotentials stands for a scattering matrix  $\mathbf{S}$  defined as

$$\begin{pmatrix} Z \\ \tilde{Z} \end{pmatrix} = \mathbf{S} \begin{pmatrix} Z' \\ \tilde{Z}' \end{pmatrix}, \quad (4)$$

which has a decomposition

$$\mathbf{S} = \begin{pmatrix} e^{i\phi} & 0 \\ 0 & e^{i\tilde{\phi}} \end{pmatrix} \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} e^{i\phi'} & 0 \\ 0 & e^{i\tilde{\phi}'} \end{pmatrix}, \quad (5)$$

Such a form of  $\mathbf{S}$  provides the conservation of flux  $|Z|^2 + |\tilde{Z}|^2 = |Z'|^2 + |\tilde{Z}'|^2$ . The coupling strength is characterized by an angle  $\alpha$  (gauge phases in matrix  $\mathbf{S}$ , as well as in  $\mathbf{M}$ , can be absorbed into  $Z$ 's). It is easy to see that the only effect of such a coupling to a loop is just a phase shift between amplitudes  $Z$  and  $Z'$ . Indeed,  $\tilde{Z}$  and  $\tilde{Z}'$  differ by a phase factor acquired by an electron traversing the loop:  $\tilde{Z}' = \tilde{Z}e^{i\varphi}$ . Then (4) yields  $Z = Z'e^{i\delta}$ , with

$$\tan \delta(E) = \frac{\sin \varphi \sin^2 \alpha}{\cos \varphi (1 + \cos^2 \alpha) - 2 \cos \alpha}. \quad (6)$$

The energy dependence of the scattering phase  $\delta$  is determined by the energy dependence of  $\varphi$ . For a small coupling the resonances occur at  $\varphi = \pm \alpha^2/2 + 2\pi p$  ( $p$  is an integer) the resonance width being also  $\alpha^2/2$ . However, these resonances are of no importance since the phase  $\delta$  can also be absorbed into the random phase on the link.

The situation is completely different when a loop occurs in the vicinity of the saddle-point so that it is coupled to the both incoming and outgoing links. It is important to note that such a loop is not coupled directly to the saddle-point because the gradient of potential there is precisely zero. The tunneling between the loop and the links occurs most favorably at a certain distance from the saddle point, as illustrated in Fig. 2b. Since the characteristic tunneling distance at the saddle-point is quite small (typically, the magnetic length), we neglect the effect of the loop on tunneling transparency at the saddle-point.

Below we demonstrate that the effect of such loops is to shift the position of delocalized state up. The crucial observation is that the saddle-point with the loops attached to the links can be viewed as some modified saddle-point (see Fig. 2b) and, thus, characterized by a matrix  $\mathbf{M}'$  with the same unitarity properties

$$\begin{pmatrix} Z'_1 \\ Z'_3 \end{pmatrix} = \mathbf{M}' \begin{pmatrix} Z'_4 \\ Z'_2 \end{pmatrix}, \quad \mathbf{M}' = \begin{pmatrix} e^{i\phi'_1} & 0 \\ 0 & e^{i\phi'_2} \end{pmatrix} \begin{pmatrix} \cosh \theta' & \sinh \theta' \\ \sinh \theta' & \cosh \theta' \end{pmatrix} \begin{pmatrix} e^{i\phi'_3} & 0 \\ 0 & e^{i\phi'_4} \end{pmatrix}, \quad (7)$$

The new parameter  $\theta'$  can be expressed via  $\theta$  and the elements of scatterig matrices  $\mathbf{S}_i$  (see Fig. 2b) using the following equations

$$\begin{pmatrix} Z'_2 \\ Z'_5 \end{pmatrix} = \mathbf{S}_2 \begin{pmatrix} Z_2 \\ Z_6 \end{pmatrix}, \quad \begin{pmatrix} Z_3 \\ Z_6 \end{pmatrix} = \mathbf{S}_3 \begin{pmatrix} Z'_3 \\ Z'_5 \end{pmatrix}, \quad \begin{pmatrix} Z'_1 \\ Z'_8 \end{pmatrix} = \mathbf{S}_1 \begin{pmatrix} Z_1 \\ Z_7 \end{pmatrix}, \quad \begin{pmatrix} Z_4 \\ Z_7 \end{pmatrix} = \mathbf{S}_4 \begin{pmatrix} Z'_4 \\ Z'_8 \end{pmatrix}, \quad (8)$$

Consider for simplicity a case when all  $\alpha_i = \alpha$  and there is only one loop (e.g.,  $\mathbf{S}_1 = \mathbf{S}_4 = \mathbf{1}$ ).

Then the solution of (1), (7) and (8) reads

$$\sinh^2 \theta' = \frac{2 \sinh^2 \theta \cos^2 \alpha (1 - \cos \varphi)}{[\sin \varphi - \sin^2 \alpha \cosh \theta \sin(\psi - \varphi)]^2 + [\cos \varphi - \cos^2 \alpha + \sin^2 \alpha \cosh \theta \cos(\psi - \varphi)]^2}. \quad (9)$$

Here  $\varphi$  [the same as in (6)] is the phase acquired by electron traversing the loop while  $\psi$  is the phase acquired on the contour  $Z_2 \rightarrow Z_3 \rightarrow Z_5$  (see Fig. 2b).

The conversion of the parameter  $\theta$  into  $\theta'$  can be viewed as an effective change in the height of the saddle-point,  $\delta V_0$ , caused by  $n = 1$  LL:

$$\delta V_0 = -\gamma \ln \left( \frac{\sinh \theta'}{\sinh \theta} \right), \quad (10)$$

The distribution function of  $\delta V_0$ , which comes from averaging over the random phases  $\varphi$  and  $\psi$ , is plotted in Fig. 3 for different values of  $\alpha$ . The narrow peaks at small  $\delta V_0$  originate from the values of  $\varphi$  that are not too close to  $2\pi p$ . Then it is easy to see from (9) that for such  $\varphi$  and  $\alpha \ll 1$  we have  $\theta' \simeq \theta$  and, consequently,  $\delta V_0$  is small. At the same time, the tails of the distribution come from the resonances  $\varphi \simeq 2\pi p$ . For such  $\varphi$  we may have both  $\theta' > \theta$  and  $\theta' < \theta$ . Important is, however, that the tails are *asymmetric*, i.e., the distribution function falls off slower towards large  $\delta V_0$ . As a result, the average  $\overline{\delta V_0}$  appears to be positive. For small  $\alpha$  this average can be easily calculated analytically:

$$\overline{\delta V_0}(\theta) = \frac{\alpha^2 \gamma}{\pi} [\sinh \theta + \arcsin(1/\cosh \theta)], \quad (11)$$

where  $\theta$  is related to  $V_0$  by Eq. (3). We see that  $\overline{\delta V_0}(\theta)$  is proportional to the width of the resonance,  $\alpha^2$ , since it is determined by resonant loops. Since  $\overline{\delta V_0}(\theta)$  is finite, the average saddle-point height,  $\overline{V_0}$ , moves from  $\overline{V_0} = 0$  to a *finite* value  $\overline{V_0} = \langle \overline{\delta V_0}(\theta) \rangle_{V_0}$ , where  $\langle \rangle_{V_0}$  stands for averaging over  $V_0$ . The values of  $V_0$ , relevant for delocalization, are of the order of  $\gamma$ , the relevant  $E - V_0$  being also of the order of  $\gamma$ , so that the relevant  $\theta$  in (11) is  $\theta \sim 1$ . This leads to the following estimate for the energy shift of the delocalized state:

$$\delta E_0 \sim \langle \overline{V_0} \rangle_\alpha \sim \overline{\alpha^2} \gamma, \quad (12)$$

where  $\overline{\alpha^2}$  is the coupling strength averaged over the loops. It can be seen that the shift is much smaller than a typical saddle-point height:  $\delta E_0/\gamma \sim \overline{\alpha^2} \ll 1$ .

Physically, the levitation (positive  $\delta E_0$ ) originates from the loops providing a direct transmission between links [19], bypassing the saddle-point. In fact, some loops cause an

opposite effect: by acquiring an appropriate phase  $\psi$  on the contour  $Z_2 \rightarrow Z_3 \rightarrow Z_5$  the electron can be better transmitted to the other side of the saddle-point. However, this effect appears to be smaller than the effect of bypassing. Thus, in order to compensate the leakage of electrons to the opposite link via the loop, the energy of the delocalized state raises up.

In the derivation of (11) we neglected the effect of the loop  $Z_7 \rightarrow Z_8$  at the other side of the saddle-point (see Fig. 2b). It is easy to see, however, that the two loops cause essentially an additive effect. Indeed, an electron, traversing the link  $Z'_1$ , can also bypass the saddle-point due to a resonant transmission to the opposite link  $Z'_4$ .

The key-point of the above consideration was that the loops can occur only in directions of the descend of a saddle-point potential (valleys), i.e., to the right and to the left from the saddle-point in Fig. 2b. This is the case when the effect of  $n = 1$  LL on  $n = 0$  delocalized state is studied. For  $n = 1$  delocalized state the situation is more complicated: the loops from  $n = 2$  LL occur in the directions to the valleys while the loops from  $n = 0$  LL occur in the directions to the hills. The latter loops cause an opposite trend, pulling the delocalized state down. The resulting shift can be presented as  $\delta E_1 \sim \gamma(\overline{\alpha_{12}^2} - \overline{\alpha_{10}^2})$ . Note that the coupling,  $\overline{\alpha_{12}^2}$ , of  $n = 1$  to  $n = 2$  LL is stronger than the coupling,  $\overline{\alpha_{10}^2}$ , of  $n = 1$  to  $n = 0$  LL due to a larger size of the  $n = 2$  wave function. Thus,  $\delta E_1$  is also positive. Another effect which we have neglected is coupling of the  $n = 0$  links via the loops corresponding to  $n = 0$  equipotentials. Since these loops occur with equal probability in *all* directions from the saddle-point, it is obvious that positive and negative contributions to  $\delta E_0$  from these loops cancel each other out.

The effective coupling constants  $\overline{\alpha_{n,n+1}^2}$  are small if the LL width is small, i.e.,  $W \ll \hbar\omega_c$ . But even if  $W \gg \hbar\omega_c$ , the coupling can be still small since it is determined by tunneling between equipotentials. A crude estimate for a tunneling amplitude in this case,  $W \gg \hbar\omega_c$ , can be obtained as follows. The spacial distance between equipotentials is of the order of  $s = \hbar\omega_c / (\frac{W}{R_c})$ ,  $R_c$  being the correlation radius of a smooth potential. This corresponds to the momentum transfer  $q = s/l^2$ ,  $l$  being the magnetic length. The coupling is efficient only if  $q < R_c^{-1}$ . Otherwise the smooth potential will not be able to provide the necessary

momentum, and the tunneling amplitude would be exponentially small. The latter condition,  $qR_c \lesssim 1$ , can be rewritten as

$$\hbar\omega_c \lesssim \left[ W \left( \frac{\hbar^2}{mR_c^2} \right) \right]^{1/2} \equiv \hbar\omega_c^*. \quad (13)$$

Eq. (13) requires that the magnetic field should not be too strong; however, the quasiclassical picture of electron motion along equipotentials still applies at  $\omega_c \sim \omega_c^*$ : for such  $\omega_c$  we have  $l/R_c \sim [(\frac{\hbar^2}{mR_c^2})/W]^{1/4} \ll 1$ , which is just the condition that the potential is smooth. On the other hand, the use of the Chalker-Coddington model is justified only if the saddle-point “discriminates” neighboring LL, i.e.,  $\hbar\omega \gtrsim \gamma$ . Remarkably, the two conditions,  $\omega_c \sim \omega_c^*$  and  $\hbar\omega_c \sim \gamma$ , coincide. Indeed, following Polyakov and Shklovskii [20], the parameter  $\gamma$  in (3) can be estimated as  $\gamma \sim \hbar(\frac{W}{mR_c^2})/\omega_c \sim \hbar\omega_c^{*2}/\omega_c$ . Thus, the Chalker-Coddington model is inadequate for  $\omega_c \ll \omega_c^*$ , when the saddle-point at  $\hbar\omega_c(n + 1/2)$  becomes transparent also for several neighboring to  $n$ th LL; for such fields one might expect a strong levitation. Our consideration applies for magnetic fields  $\omega_c > \omega_c^*$ , when the levitation is weak.

Let us briefly discuss the dependence of the coupling on LL number  $n$ . This dependence is determined by the overlap of  $n$ th and  $(n + 1)$ th Landau wave-functions shifted in space by a distance  $s$ . This overlap is known to be proportional to  $e^{-u/2} L_n^1(u) [u/(n + 1)]^{1/2}$ , where  $u = s^2/2l^2$  and  $L_n^1$  is the Laguerre polynomial. If the ratio  $\omega_c/\omega_c^* = (qR_c)^{1/2} = (R_c s/l^2)^{1/2}$  is not too large (the coupling is not exponentially small) we have  $s < l$ . Then it is easy to see that  $\overline{\alpha_{n,n+1}^2} \propto [L_n^1(0)]^2/n \propto n$ . As a result, the levitation  $\delta E_n$ , which is proportional to  $\overline{\alpha_{n,n+1}^2} - \overline{\alpha_{n,n-1}^2}$ , does not depend on  $n$ .

In conclusion, we have traced how the LL mixing gives rise to a shift of the energy of the delocalized state up from the position  $\hbar\omega_c(n + 1/2)$ . In a smooth potential the shift may be relatively small even when the peaks in the density of states corresponding to different LL are not well-resolved [it can be seen from (13) that  $\hbar\omega_c^*/W = [(\frac{\hbar^2}{mR_c^2})/W]^{1/2} \ll 1$ ]. Surprisingly, the simulation [15] for the *white-noise* random potential, where one might expect much stronger coupling between LL, indicated that for  $\hbar\omega_c \simeq W$  the levitation is still very weak.

The picture considered above was a single-particle picture of delocalization. In other

words, we have completely neglected the effects of screening caused by electron-electron interactions. At the same time, it is known that in a smooth potential the electron-electron interactions affect drastically the distribution of electrons within the plane. In fact, when the potential is very smooth the equipotentials, we have dealt with, are separated by incompressible strips [21]. However, as it was discussed, a noticeable levitation occurs only when the field is not too strong, so that the distance between equipotentials is less than the magnetic length. Clearly, in this case the macroscopic description of [21] does not apply and a single-particle picture should be valid.

### ACKNOWLEDGMENTS

Authors acknowledge the discussions with D. E. Khmelnitskii and D. G. Polyakov on subjects related to this work.

## REFERENCES

- [1] B. I. Halperin, Rhys. Rev. B **25**, 2185 (1982).
- [2] H. Levine, S. B. Libby, and A. M. M. Pruisken, Rhys. Rev. Lett. **51**, 1915 (1983).
- [3] D. E. Khmelnitskii, Pis'ma Zh. Eksp. Teor. Fiz. **38**, 454 (1983) [JETP Lett. **38**, 552 (1983)].
- [4] H. Aoki and T. Ando, Rhys. Rev. Lett. **54**, 831 (1985); see also H. Kamimura and H. Aoki, *The Physics of Interacting Electrons in Disordered Systems* (Clarendon Press, Oxford, 1989) and references therein.
- [5] B. Huckenstein and B. Kramer, Rhys. Rev. Lett. **64**, 2304 (1990); see also recent reviews M. Janssen, O. Viehweger, U. Fastenrath, and J. Hajdu, *Introduction to the Theory of the Integer Quantum Hall Effect* (VCH, Weinheim, 1994), and B. Huckenstein, MPI preprint, 1995 (cond-mat/9501106).
- [6] Y. Huo and R. N. Bhatt, Rhys. Rev. Lett. **68**, 1375 (1992).
- [7] D. E. Khmelnitskii, Phys. Lett. **106A**, 182 (1984).
- [8] R. B. Laughlin, Rhys. Rev. Lett. **52**, 2304 (1984).
- [9] S. Kivelson, D.-H. Lee, and S.-C. Zhang, Phys. Rev. B **46**, 2223 (1992).
- [10] H. W. Jiang, C. E. Johnson, K. L. Wang, and S. T. Hannahs, Rhys. Rev. Lett. **71**, 1439 (1993).
- [11] C. E. Johnson and H. W. Jiang, Rhys. Rev. B **48**, 2823 (1993).
- [12] T. Wang, K. P. Clark, G. F. Spencer, A. M. Mack, and W. P. Kirk, Phys. Rev. Lett. **72**, 1439 (1994).
- [13] R. J. F. Hughes, J. T. Nicholls, J. E. F. Frost, E. H. Linfield, M. Pepper, C. J. B. Ford, D. A. Ritchie, G. A. C. Johns, E. Kogan, and M. Kaveh, J. Phys.: Condens. Matter **6**,


- 4763 (1994).
- [14] I. Glozman, C. E. Johnson, and H. W. Jiang, Rhys. Rev. Lett. **74**, 594 (1995).
  - [15] T. Ando, J. Rhys. Soc. Jpn. **53**, 3126 (1984)
  - [16] J. T. Chalker and P. D. Coddington, J. Phys. C **21**, 2665 (1988).
  - [17] D.-H. Lee, Z. Wang, and S. Kivelson, Phys. Rev. Lett. **70**, 4130 (1993).
  - [18] H. A. Fertig and B. I. Halperin, Phys. Rev. B **36**, 7969 (1987).
  - [19] A similar process of a resonant tunneling between the edges of a Quantum Hall sample was considered in J. K. Jain and S. A. Kivelson, Phys. Rev. Lett. **60**, 1542 (1988).
  - [20] D. G. Polyakov and B. I. Shklovskii, Phys. Rev. Lett. **73**, 1150 (1994); *ibid* **74**, 150 (1995)
  - [21] A. L. Efros, Solid State Comm., **67**, 1019 (1988); D. B. Chklovskii, B. I. Shklovskii, and L. I. Glazman, Phys. Rev. B **46**, 4026 (1992); N. R. Cooper and J. T. Chalker, *ibid* **48**, 4530 (1993).


## FIGURES

FIG. 1. Schematic picture of the levitation of delocalized states according to [7,8]. The straight lines correspond to  $n = 0$  and  $n = 1$  Landau levels.


FIG. 2. Sketch of the saddle-point. The full lines represent equipotentials corresponding to  $n = 0$  LL. The dashed circles are equipotentials of  $n = 1$  LL. (a) Scattering of an electron by isolated loop (**S** stands for the scattering matrix). (b) Passage of electron through the saddle-point in the presence of loops connecting the links.

FIG. 3. The distribution function of  $\delta V_0$  is shown at  $E = V_0$  ( $\theta = \theta_c$ ) for  $\alpha^2 = 0.1$  (solid line),  $\alpha^2 = 0.2$  (long-dashed line), and  $\alpha^2 = 0.3$  (dashed line).





(a)



(b)

